

DARLEX[®]

Insecticide

Specimen Label

Darlex[®] Insecticide controls darkling beetles (also known as lesser mealworms) and hide beetles in poultry housing. This product can be used as a crack and crevice, general surface or spot treatment applied to walls, floors and support beams of poultry (broiler, turkey, caged egg layer, breeder) housing.

- For use in, and along the outside perimeter of poultry houses to control darkling beetles and hide beetles
- Controls adult and larvae darkling beetles
- For broadcast and band applications
- Specially formulated for the poultry industry
- Easy to mix and spray



ACTIVE INGREDIENT:

| | |
|-------------------------|---------|
| *Clothianidin | 23.00% |
| OTHER INGREDIENTS | 77.00% |
| | 100.00% |

*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine
Contains 2.13 lbs. active ingredient per gallon.

Darlex is a registered trademark of McLaughlin Gormley King Company.

PRECAUCIÓN AL CONSUMIDOR: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

NOT REGISTERED FOR USE, SALE OR DISTRIBUTION IN THE STATE OF NEW YORK.

- Do not apply this product in any way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.
- Poultry producers should use a range of management practices to reduce the possibility of insecticide resistance and to help minimize the buildup and carryover of populations of darkling beetles and other listed insect pests. Contact your local Cooperative Extension office for advice concerning the use of this product and appropriate resistance management strategies.
- **Shake well before using.**
- Darlex Insecticide controls darkling beetles (also known as lesser mealworms) and hide beetles in poultry housing. This product can be used as a crack and crevice, general surface, or spot treatment applied to walls, floors and support beams of poultry (broiler, turkey, caged egg layer, breeder) housing.

- **Do not apply when birds are present** and do not expose poultry to direct spray. Exposed water lines and feed troughs must be covered or removed from the area before treatment begins. Once completed, allow surfaces to dry completely before restocking. Do not apply as a fog or space spray. Fresh bedding material must be put into the house prior to restocking.
- **Read and follow ALL label directions when applying this product.**
- Do not use Darlex Insecticide at less than the label rate. By using less than the specified rate, insects are exposed to sub-lethal doses, which may not kill the insect, but can increase the development of resistance.

When To Apply

Apply after de-caking, clean-out and sanitation procedures are complete, but prior to new flock placement. It may be advisable to treat houses before removing litter to prevent possible infestation of nearby residences.

Mixing Instructions

Mix 4 ounces in ½–2 gallons of water to treat 1,000 sq. ft. Be sure to accurately measure the actual surface area to be treated in order to determine the quantity of Darlex Insecticide required for the application. To prepare the diluted spray, fill the sprayer tank with half the total water to be used, add the appropriate amount of Darlex Insecticide, then add the remaining water while agitating or mixing.

To apply 4 oz. Darlex Insecticide per 1,000 sq. ft.

| Sq. Ft. of Area to Be Treated | Fluid Ounces of Darlex Insecticide Needed | Total Volume of Water |
|-------------------------------|---|-----------------------|
| 500 | 2 fl. oz. | 1/4–1 gallon |
| 5,000 | 20 fl. oz. | 2.5–10 gallons |
| 10,000 | 40 fl. oz. | 5–20 gallons |
| 15,000 | 60 fl. oz. | 7.5–30 gallons |
| 20,000 | 80 fl. oz. | 10–40 gallons |

Make certain that sufficient agitation is maintained throughout the application to ensure a uniform spray. Prepare a fresh spray mixture before each treatment.

NOTE: Adding disinfectants to insecticide tank mixes may affect insecticidal activity.

APPLICATION METHODS

Brand Application: Darkling beetles often crowd under feeders, water lines or along perimeter walls, lying buried in the litter. Under these circumstances, it may be sufficient to treat the primary areas of beetle infestation within a poultry (broiler, turkey, caged egg layer, breeder) house. Beetles can also infest structures behind insulation in walls; Darlex Insecticide can be applied as a spot treatment inside wall voids or as a crack and crevice application to target these infestations.

For band application, apply diluted product to a 3–5 foot wide swath of floor area, making sure to get under all the water and/or feed lines in the house. Continue applying a 3–5 foot swath along the inside perimeter of the side and end walls, including the entire surface of all footings and 1–2 feet up onto the walls, support beams and wood surfaces above the foundation.

Interior Broadcast Application: Whenever an infestation is severe, it may be necessary to treat the entire house by applying the diluted product as a broadcast spray. Spray Darlex Insecticide on the entire floor, paying particular attention to areas beneath feed and water lines, as well as to the lower sections of walls up to 1 foot on wood surfaces above the foundation. Fresh bedding must be put into the house prior to restocking.

For Best Results: In houses with support beams, treat around the base of each support post and 1 foot up each post. Apply diluted spray to cracks and crevices around insulation where beetles have been seen or can find harborage. To prevent movement of beetle infestations to houses and nearby residences, it may also be desirable to treat manure before it is relocated from the poultry house.

Use Darlex Insecticide as part of an integrated pest management program to extend the efficacy of insect control products and to help minimize the buildup and carryover of resistant insect populations. Darkling beetles have the ability to develop resistance to insecticides. When one class of chemistry is used year after year, the likelihood that insects will develop resistance and become more difficult to control, greatly increases. This product contains clothianidin, a neonicotinoid insecticide which should be used in an insecticide rotation program with other classes of insecticides, such as pyrethroids, organophosphates or spinosyns, to minimize or prevent the development of resistance and to maintain its effectiveness in controlling darkling beetles and other listed insects.

In order to avoid re-infestation after cleanout, apply targeted applications to the exterior of the building perimeter, foundation, doors and windows, and other places where pests may enter the building. Do not allow this product to contact plants in bloom if bees are foraging the treatment area.

Darlex Insecticide may be used to control crickets and nuisance ants in, and along the outside perimeter of animal housing. Mix 1/2–1 ounce of Darlex Insecticide per gallon of water. Apply as a spot treatment, banded application or crack and crevice application at the base of the structure foundation or at entry points such as doors, vents, eaves or utility penetrations. In building interiors, apply only as a crack and crevice or void application. If desired, spot or drench treatments may be made to soil around the structure exterior where ants nest. Above ground ant nests in wood posts, decks, or fences, may be treated by spraying the cavity where ants are nesting or the wood surface.

Not for use in control of red imported fire ants, native fire ants, pharaoh ants, harvester ants or carpenter ants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a dry place away from excessive heat. Do not store near food or feed. Store in original container only. To close package, replace and tighten cap to form an airtight seal.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label instructions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with solvent used to dilute the product for use and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride, or viton.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply when weather conditions favor drift from treated areas. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

DARLEX[®]
Insecticide

EPA Reg. No. 1021-2771
EPA Est. 1021-MN-2
Rev. 0815-0915

MGK[®]

Manufactured by MGK[®].
©2015 MGK. All rights reserved.
8810 Tenth Avenue North, Minneapolis, MN 55427

2.5M-1111