

## TETROXY® HCA™ For Poultry and Swine

Soluble Powders  
(Oxytetracycline HCl) Antibiotic

### Active Ingredient

Oxytetracycline Hydrochloride

### Indications

For use in the drinking water of poultry and swine as a control of specific diseases (see back page) susceptible to oxytetracycline.

### Benefits

- **Versatile**—Antibiotic treatment for chickens, turkeys and swine
- **Safe**—Zero withdrawal time before slaughter
- **Convenient, accurate, economical**—Full range of packaging options for the way producers work today
- **Soluble**—Readily dissolves in water
- **Easy to use and store**—Pre-measured packets packaged in space-saving plastic pails and economical 5 lb bulk pail
- **FDA approved**

### Administration\*

Mix fresh solutions daily. Use as sole source of drinking water. Do not mix product directly with milk or milk replacers. Administer one hour before or two hours after feeding milk or milk replacers.

Consult a poultry veterinarian or diagnostic laboratory. As a generalization, 200 chickens will drink one gallon of water per day per each week of age. Turkeys will consume twice that amount. Administer 7 to 14 days to chickens and turkeys and up to five days to swine.



List No.	UNIT PACKAGE
1TET012	TETROXY HCA-280 9.87 oz. packets (40 packets per pail)
1TET013	TETROXY HCA-560 19.74 oz. packets (25 packets per pail)
1TET014	TETROXY HCA-1400 3.09 lb. packets (10 packets per pail)
1TET019	TETROXY HCA-2270 5 lb. packets (6 packets per pail)
1TET011	TETROXY HCA-2270 5 lb. bulk pail (6 pails per case)

\*See inside for complete dosing information.

# Technical Information

## TETROXY® HCA-1400

### ACTIVE INGREDIENT(S)

Each packet contains 512 g oxytetracycline HCl and will make:  
 2560 gal. (9690 L) containing 200 mg oxytetracycline HCl per gal.  
 1280 gal. (4845 L) containing 400 mg oxytetracycline HCl per gal.  
 640 gal. (2422 L) containing 800 mg oxytetracycline HCl per gal.  
 3.09 lbs. (1.40 kg) will treat 51,200 pounds of swine.

### PRESENTATION

3.09 lbs. (1400 g) packets. 10 packets per resealable pail.

## MIXING INSTRUCTIONS FOR WATER MEDICATION CHICKENS-TURKEYS-SWINE

**DOSAGE AND INDICATIONS FOR USE IN WATER PROPORTIONERS:** Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	Disease	Treatment Level	Gallons Stock/ Packet
CHICKENS	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-800 mg	10-5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400-800 mg	10-5
TURKEYS	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	GROWING TURKEYS: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
SWINE	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

TAKE TIME



OBSERVE LABEL  
 DIRECTIONS

# Technical Information

## **TETROXY® HCA™** (Oxytetracycline HCl) ANADA 200-144, Approved by FDA

### INDICATIONS

For use in the drinking water of poultry and swine as a control of specific diseases susceptible to oxytetracycline.

### DOSAGE & ADMINISTRATION

Mixing instructions for water medication: Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon. See chart below.

### PRECAUTIONS

Use as sole source of Oxytetracycline. Not to be used for more than 14 consecutive days in chickens and turkeys or 5 consecutive days in swine.

### WARNING

Do not feed to birds producing eggs for human consumption. Discontinue treatment of chickens and turkeys and swine 0 days prior to slaughter. Consult material safety data sheet prior to use.

## **TETROXY® HCA™-280**

### ACTIVE INGREDIENT(S)

Each packet contains 102.4 g oxytetracycline HCl and will make: 512 gal. (1938 L) containing 200 mg oxytetracycline HCl per gal. 128 gal. (484 L) containing 800 mg oxytetracycline HCl per gal.

### PRESENTATION

9.87 oz. (280 g) packets. 40 packets per resealable pail.

## **TETROXY® HCA™-560**

### ACTIVE INGREDIENT(S)

Each packet contains 204.8 g oxytetracycline HCl and will make: 1024 gal. (3876 L) containing 200 mg oxytetracycline HCl per gal. 512 gal. (1938 L) containing 400 mg oxytetracycline HCl per gal. 256 gal. (969 L) containing 800 mg oxytetracycline HCl per gal. One packet will treat 20,480 pounds of swine.

### PRESENTATION

19.74 oz. (560 g) packets. 25 packets per resealable pail.

## MIXING INSTRUCTIONS FOR WATER MEDICATION CHICKENS-TURKEYS-SWINE

**DOSAGE AND INDICATIONS FOR USE IN WATER PROPORTIONERS:** Add the following amount **TETROXY® HCA™-280** to **One Gallon** of stock solution or **TETROXY® HCA™-560** to **Two Gallons** of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

	Disease	Treatment Level	280/gal. or 560/2 gal. Stock Solution
<b>CHICKENS</b>	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	1/4-1/2 (34-68 g)
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-800 mg	1/2-1 (68-135.5 g)
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400-800 mg	1/2-1 (68-135.5 g)
<b>TURKEYS</b>	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	1/4-1/2 (34-68 g)
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	1/2 (68 g)
	GROWING TURKEYS: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
<b>SWINE</b>	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

# Technical Information

## TETROXY® HCA-2270

### ACTIVE INGREDIENT(S)

Each packet contains 830.2 g oxytetracycline HCl and will make:  
4151 gal. (15712 L) containing 200 mg oxytetracycline HCl per gal.  
2075 gal. (7854 L) containing 400 mg oxytetracycline HCl per gal.  
1038 gal. (3929 L) containing 800 mg oxytetracycline HCl per gal.  
5 lbs. (2.27 kg) will treat 83,017 pounds of swine.

### PRESENTATION

5 lbs. (2270 g) packets; 6 packets per resealable pail and 5 lbs. resealable bulk pail.

## MIXING INSTRUCTIONS FOR WATER MEDICATION

### CHICKENS-TURKEYS-SWINE

**DOSAGE AND INDICATIONS FOR USE IN WATER PROPORTIONERS:** Add the following amount to one gallon of stock solution when proportioner is set to meter at the rate of ounce per gallon.

	Disease	Treatment Level	Quantity/Gallon Stock Solution
CHICKENS	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	70-140 g
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-800 mg	140-280 g
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400-800 mg	140-280 g
TURKEYS	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	70-140 g
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	140 g
	GROWING TURKEYS: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
SWINE	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

Manufactured by:



Bimeda, Inc.  
Riverside, MO 64150

TETROXY is a registered trademark of Bimeda, Inc. HCA is a trademark of Bimeda, Inc.