

AmproMed™ P FOR POULTRY

(amprolium)

20% Soluble Powder/Coccidiostat

INDICATIONS

AmproMed™ P FOR POULTRY is intended for the treatment of coccidiosis in growing chickens, turkeys and laying hens.

BENEFITS

- **Effective:** breaks up the coccidia life cycle
- **Versatile:** labeled for the treatment of coccidiosis in growing chickens, turkeys and laying hens
- **Easy To Use:** suitable for use in automatic water proportioners
- **Convenient:** treats large numbers of birds through the drinking water
- **Economical:** a single 10 oz pack makes 128 gallons of medicated drinking water at 0.012% level
- **Palatable:** accepted by poultry in the drinking water
- **Safe:** approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1AMP022	10 oz (283.5 g) pouch	24 per pail

See reverse side for Administration and Dosage.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com



AmproMed™ P FOR POULTRY

(amprolium)

20% Soluble Powder/Coccidiostat

ANADA 200-488, Approved by FDA

NOT FOR HUMAN USE

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN

Water-Soluble Treatment for Coccidiosis

Contents of each packet will treat 128 gallons at 0.012% level

Restricted Drug (California) – Use Only as Directed

INDICATIONS:

AmproMed™ P FOR POULTRY is intended for the treatment of coccidiosis in growing chickens, turkeys and laying hens.

USE DIRECTIONS:

1 oz = 3 1/2 measuring tablespoonfuls

PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:

For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

To prepare dosage levels of:	Dissolve AmproMed™ P FOR POULTRY in one gallon of water. (Makes 1 gallon of stock solution.)
0.024%	2 bags (20 oz) in 1 gallon
0.012%	1 bag (10 oz) in 1 gallon
0.006%	1/2 bag (5 oz) in 1 gallon

To prepare 50 Gallons of Medicated Water:

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Dissolve 8 ounces of AmproMed™ P FOR POULTRY in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use 4 ounces of AmproMed™ P FOR POULTRY.
0.006%	Follow same directions as above but use 2 ounces of AmproMed™ P FOR POULTRY.

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

USE DIRECTIONS:

Give amprolium at the 0.012% level (10 oz AmproMed™ P FOR POULTRY per 128 Gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level.) Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as sole source of amprolium.

CAUTION:

If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

STORAGE:

Store at 20°C-25°C (68°F-77°F), excursions permitted between 15°C-30°C (59°F-86°F).

To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

