

HUVEPHARMA, INC.

525 WESTPARK DR., SUITE 230, PEACHTREE CITY, GA, 30269

Telephone: 770-486-7212

Toll-Free: 877-994-4883

Fax: 770-486-7217

Website: www.huvepharma.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

AMPROL® 128 20% SOLUBLE POWDER

Huvepharma

(amprolium)

Coccidiostat

Water-Soluble Treatment for Coccidiosis

Contents of each packet will treat 128 gallons at 0.012% level

PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS

For Automatic Water Proportioners

For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

To prepare dosage levels of:	Dissolve AMPROL 128 (amprolium) 20% Soluble Powder in one gallon of water. (Makes 1 gallon of stock solution.)
0.024%	2 bags (20 oz) in 1 gallon
0.012%	1 bag (10 oz) in 1 gallon
0.006%	1/2 bag (5 oz) in 1 gallon

To Prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Dissolve 8 ounces of AMPROL 128 20% Soluble Powder in about 5 gallons of water in a 50-gallon medication barrel. Stir, then add water to the 50-gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use 4 ounces of AMPROL 128 20% Soluble Powder.
0.006%	Follow same directions as above but use 2 ounces of AMPROL 128 20% Soluble Powder.

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

FOR ANIMAL USE ONLY

AMPROL® 128 (amprolium) 20% Soluble Powder is intended for the treatment of coccidiosis in growing chickens, turkeys and laying hens.

CAUTION: If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

NOT FOR HUMAN USE.

Restricted Drug (California) - Use Only as Directed.

Keep this and all drugs out of the reach of children.

STORAGE: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C (104°F).

USE DIRECTIONS: Give amprolium at the 0.012% level (10 oz AMPROL 128 20% Soluble Powder per 128 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level.) Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

NADA 33-165, Approved by the FDA

Distributed by: Huvepharma, Inc., Peachtree City, GA 30269

For questions, please call 1-877-4AMPROL

AMPROL® is a registered trademark of Huvepharma, Inc.

Net Contents:	Product #	
10 oz. (283.5 grams)	970915	1075-1816-02
		Rev. 07-2008

NAC No.: 13090080