

FLEAS: General Volumetric Treatment: Hold aerosol 36 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor areas are contacted. Change litter before application. Keep area closed for 15 minutes. Remove animals before spraying. Open and ventilate before reoccupying. Repeat treatment after 7 days or as necessary.

KENNELS AND CATTERIES: Remove animals before spraying. Ventilate after treatment is completed.

OUTDOOR GROUND APPLICATIONS: Flies, gnats and mosquitoes in open areas near buildings and in campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60-80 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION

Harmful if absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **HOT LINE NUMBER –** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985)878-6751, Monday through Friday, 8:00am to 4:30pm CST. After 4:30pm call your poison control center. For product information call (203)597-1812, Monday-Friday, 8:00am to 5:00pm.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame. **DISPOSAL:** Do not Puncture or Incinerate!

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or

1-800-CLEANUP for disposal instructions.



For Additional Products and Information Visit
Us At www.Country-Vet.com

Manufactured by
WATERBURY COMPANIES, INC.

P.O. Box 1812 • Waterbury, CT 06722
985-878-6751 • Made in U.S.A.



INSTANT KNOCKDOWN

CV-80D

FARM & DAIRY
for Flying Insect Control

0.50% Pyrethrins
for Quick KILL

KILLS & REPELS:

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Gnats, Cockroaches, Fleas, Asian Beetles, Barn Flies, Deer Flies, Stable Flies, Horn Flies, Horse Flies, House Flies, Face Flies, Lice and Cluster Flies

FOR USE IN:

Beef Cattle Operations, Dairy Farms, (including Milk House, Milk Parlor, Loafing Sheds and Holding Lot), Hog Operations, Kennels, Barns, Stables, Farms, Animal Quarters, Milkrooms and Poultry Houses



CONTAINS NO CFCs or OTHER OZONE DEPLETING SUBSTANCES.
FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.50%
*Piperonyl Butoxide, TECHNICAL.....	4.00%
INERT INGREDIENTS.....	95.50%
	100.00%

*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% related compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements.
EPA Reg. No. 9444-210 EPA Est. No. 9444-LA-1

NET WT 18.5 oz (524 g)

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Note: When using this product in an enclosed area, such as a milkroom, barn or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milkroom: Close all windows and doors. Direct spray upwards, spraying for 1-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk before spraying to prevent contamination from spray and dead or falling insects.

ANIMAL USE: (Do not spray sick animals): Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at a rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over back, allowing 1 second per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over back of hogs and spray for 1-2 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat the application daily or as necessary.

POULTRY OPERATIONS: Use only in empty poultry houses (when birds are not present). In poultry houses, close all windows and doors. Spray roosts, walls, nests and cages thoroughly. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Treat daily or as necessary.

MUD DAUBER AND WASP NEST: Spray mist directly on insects and their nests from approximately 2 ft. distance. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

Part #34-8318CV

