

Clinafarm® EC
 Emulsifiable Concentrate

For control of
Aspergillus fumigatus
 contamination in
 poultry hatchery
 equipment



ACTIVE INGREDIENT:
 Imazalil (enilconazole): 1-(2-(2,4-Dichlorophenyl)-
 2-(2-propenyloxy)ethyl)-1H-imidazole 13.8% w/w
INERT INGREDIENTS 86.2%
 100.0% w/w

KEEP OUT OF REACH OF CHILDREN
DANGER

See back Panel for Additional Precautionary Statements

FIRST AID

If in Eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice. **If Swallowed** • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. • Promptly drink either large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. **If on Skin or Clothing** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. **If Inhaled** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling the poison control center or doctor or going for treatment. You may also contact the Rocky Mountain Poison Center at 1-303-595-4869 for emergency medical treatment information.

Net Contents: 4 - 750 mL (1.58 U.S. Pints) Bottles

Schering-Plough Animal Health
 Schering-Plough Animal Health Corp., Union, NJ 07083
 Copyright © 1998, 2003, Schering-Plough Animal Health Corp. All rights reserved.

A product of Janssen Research
 EPA Reg. No. 773-55
 EPA Est. No. 6175-LA-1

27079300 Rev. 4/03

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER

Corrosive, causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if absorbed through skin or inhaled. Avoid breathing spray mist and contact with skin. Do not get in eyes, wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For the control of *Aspergillus* organisms in cleaned poultry and turkey hatchery equipment prior to introduction of the eggs. Equipment would include empty hatcher cabinets, setters, coolers, storerooms, handling equipment, etc. CLINAFARM® EC will dramatically reduce the levels of infectious organisms and spores in treated areas. To make 1 gallon of 1:100 dilution, add 1.25 fl. oz. (36 mL) of CLINAFARM® EC to one gallon of clean water to provide an antifungal concentration of 0.15% active ingredient. Do not use product undiluted. To assure proper efficacy, the area or equipment to be treated should be clean and closed off from outside air currents before treatment. Ventilation ducts may be treated but fans must be shut off. Fog or spray area to be treated at the rate of 1/2 fl. oz. diluted product per 150 cu. ft. (1 mL per 10 cu. ft.). Fog or spray to ensure good contact with surface areas. Do not re-enter treatment areas for two hours. CLINAFARM® EC is compatible with other commonly used disinfectants such as quaternary ammonium compounds, glutaraldehyde, formaldehyde, and phenolic compounds. **PHYSICAL/CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store at room temperature. **PESTICIDE DISPOSAL:** Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CLINAFARM is a registered trademark of Johnson & Johnson CF-15

TAKE TIME



OBSERVE LABEL
 DIRECTIONS



0 1 3 8 - 5 2 0 9 - 0 1

PACKAGING COMPONENT APPROVAL

SUBMISSION # **1A**

EXAMINE BLACK & WHITE COPY AND ACCOMPANYING COLOR BREAK.

	APPROVED	NOT APPROVED	DATE
DESIGNER _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
COPY EDITOR _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
MEDICAL _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
TRADEMARKS _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
PATENTS _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
PACKAGING _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
SPECIFICATIONS _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
MARKETING _____	<input type="checkbox"/>	<input type="checkbox"/>	_____
REGULATORY _____	<input type="checkbox"/>	<input type="checkbox"/>	_____

ART DUE DATE: _____ INV. LOCATION: _____

SQA:
 APPROVAL VERIFIED: _____ DATE: _____

PANTONE COLORS

PMS 201	
PMS 422	
BLACK	