

Gallimycin® PFC

(Erythromycin) POULTRY FORMULA CONCENTRATED
WATER SOLUBLE • ANTI-INFECTIVE

NADA 035-157, Approved by FDA



INDICATIONS

IN BROILERS AND REPLACEMENT CHICKENS: CHRONIC RESPIRATORY DISEASE: As an aid in the control of Chronic Respiratory Disease due to *Mycoplasma gallisepticum* susceptible to erythromycin.

IN REPLACEMENT CHICKENS AND CHICKEN BREEDERS: INFECTIOUS CORYZA: As an aid in the control of Infectious Coryza due to *Haemophilus gallinarum* susceptible to erythromycin.

IN GROWING TURKEYS: BLUECOMB: As an aid in the control of Bluecomb (Non-specific Infectious Enteritis) caused by organisms susceptible to erythromycin.

See reverse side for Administration and Dosage.

BENEFITS

- **Effective** – contains the fast-acting antibiotic, erythromycin
- **Versatile** – can be used in broilers, replacement chickens and growing turkeys
- **Short withdrawal** – discontinue use one (1) day before slaughter
- **Economical** – one packet treats 130 gallons of drinking water
- **Convenient** – available in pre-measured packets for easy mixing

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
16AL001	250 gm	30

Gallimycin® PFC

(Erythromycin) POULTRY FORMULA CONCENTRATED

WATER SOLUBLE • ANTI-INFECTIVE

NADA 035-157, Approved by FDA

Each 250 g package contains 65 g erythromycin phosphate (Equivalent to 57.8 g erythromycin).

INDICATIONS:

IN BROILERS AND REPLACEMENT CHICKENS: CHRONIC RESPIRATORY DISEASE: As an aid in the control of Chronic Respiratory Disease due to *Mycoplasma gallisepticum* susceptible to erythromycin.

IN REPLACEMENT CHICKENS AND CHICKEN BREEDERS: INFECTIOUS CORYZA: As an aid in the control of Infectious Coryza due to *Haemophilus gallinarum* susceptible to erythromycin.

IN GROWING TURKEYS

BLUECOMB: As an aid in the control of Bluecomb (Non-specific Infectious Enteritis) caused by organisms susceptible to erythromycin.

DIRECTIONS FOR USE:

To assure effectiveness, treated birds must consume enough medicated water to provide a therapeutic dosage.

INDICATIONS:	Dosage of Erythromycin Phosphate
Chronic Respiratory Disease	1/2 g/gal of drinking water for 5 days
Infectious Coryza	1/2 g/gal of drinking water for 7 days
Blucomb (Non-specific Infectious Enteritis)	1/2 g/gal of drinking water for 7 days

MIXING INSTRUCTIONS FOR DRINKING WATER:

Add one (1) package (250 g) per 130 gallons and mix thoroughly. For proportioners which meter 1 oz/gal, add one (1) package (250 g) to each gallon of stock solution.

In chickens, this dosage provides 5.0 to 17.8 mg/lb and, in turkeys, 3.1 to 34.2 mg/lb body weight per bird per day of erythromycin activity depending upon age, class of chicken (or turkey), feed conversion, environmental temperature, and relative humidity during medication period.

Restricted Drug (California) - Use Only as Directed

WARNING:

Do **not** use in chickens or turkeys producing eggs for human consumption.

Do **not** use in replacement pullets over 16 weeks of age. Withdraw one (1) day before slaughter.

Important: Solutions older than 3 days should not be used.

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN



Gallimycin is a Registered Trademark of Bimeda, Inc.