

HUVEPHARMA, INC.

525 WESTPARK DR., SUITE 230, PEACHTREE CITY, GA, 30269

Telephone: 770-486-7212

Toll-Free: 877-994-4883

Fax: 770-486-7217

Website: www.huvepharma.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

OXYTET® SOLUBLE



Huvepharma

(oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

The 280 g packet contains 102.4 g of oxytetracycline HCl and will make:
512 gallons (1936 liters) containing 200 mg oxytetracycline HCl per gallon
128 gallons (484 liters) containing 800 mg oxytetracycline HCl per gallon
The 280 g packet will treat 10,240 lbs of swine.

The 280 g packet will treat 4,096 lbs of turkey at 25 mg/lb body weight.

The 1400 g packet contains 512 g of oxytetracycline HCl and will make:
2560 gallons (9690 liters) containing 200 mg oxytetracycline HCl per gallon
1280 gallons (4845 liters) containing 400 mg oxytetracycline HCl per gallon
640 gallons (2422 liters) containing 800 mg oxytetracycline HCl per gallon
The 1400 g packet will treat 51,200 lbs of swine.

The 1400 g packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

The 1772 g packet contains 648 g of oxytetracycline HCl and will make:
3240 gallons (12,265 liters) containing 200 mg oxytetracycline HCl per gallon
1620 gallons (6,132 liters) containing 400 mg oxytetracycline HCl per gallon
810 gallons (3,066 liters) containing 800 mg oxytetracycline HCl per gallon
The 1772 g packet will treat 64,800 lbs of swine.

The 1772 g packet will treat 25,920 lbs of turkey at 25 mg/lbs body weight.

MIXING DIRECTIONS FOR WATER MEDICATION

CHICKENS - TURKEYS - SWINE

Dosage and indications for use in water proportioners - Add the following amount to 1 gallon of stock solution when proportioner is set to meter at the rate of 1 oz per gallon.

Disease	Treatment Level	Gallons Stock/Pack		
		280 g packet	1400 g packet	1772 g packet
CHICKENS				
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	1/4 - 1/2 (70 - 140 g)	20 - 10	25.3 - 12.6
Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma</i>	400 - 800 mg	1/2 - 1 (140 - 280 g)	10 - 5	12.6 - 6.3

<i>gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.				
Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	1/2 - 1 (140 - 280 g)	10 - 5	12.6 - 6.3
TURKEYS				
Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	1/4 - 1/2 (70 - 140 g)	20 - 10	25.3 - 12.6
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	1/2 (140 g)	10	12.6
GROWING TURKEYS - Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption		
SWINE				
FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE - Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline. FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption		

USE DIRECTIONS: Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING - Do not feed to birds producing eggs for human consumption.

RECOMMENDED PACKET STORAGE CONDITIONS:

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.

NADA #130-435, Approved by FDA

Manufactured for: Huvepharma, Inc., 525 Westpark Drive, Suite 230, Peachtree City, GA 30269

HUVEPHARMA and OxyTet are registered trademarks of Huvepharma EOOD

Net Weight:	
9.87 oz (280 g)	13925000

3.09 lbs (1400 g)	13924800
3.91 lbs (1772 g)	13925200

CPN: 1309051.0