



Oxytet[®] Soluble

(oxytetracycline hcl)

DESCRIPTION

Oxytet[®] Soluble (oxytetracycline hcl soluble powder) is a water-soluble antibiotic for use in drinking water for chickens, turkeys and swine.

PACKAGING

Available in three convenient sizes:

- 50 x 280g packs/pail
- 10 x 1400g packs/pail
- 6 x 1772g packs/pail



INDICATIONS

Chickens: Control of infectious synovitis caused by *Mycoplasma synoviae*, susceptible to oxytetracycline. Control of chronic respiratory disease (CRD) and air sac infections caused by *Mycoplasma gallisepticum* and *Escherichia coli*, susceptible to oxytetracycline. Control of fowl cholera caused by *Pasteurella multocida*, susceptible to oxytetracycline.

Turkeys: Control of hexamitiasis caused by *Hexamita meleagridis*, susceptible to oxytetracycline. Control of infectious synovitis caused by *Mycoplasma synoviae*, susceptible to oxytetracycline. **Growing Turkeys** - Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.

Swine: For the control and treatment of the following diseases: Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*, susceptible to oxytetracycline and bacterial pneumonia caused by *Pasteurella multocida*, susceptible to oxytetracycline.

For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by *Leptospira pomona*, susceptible to oxytetracycline.

DOSAGE

See label (back) for full mixing directions. Administration directions and packet contents are dependent upon pack size.

MIXING DIRECTIONS

Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. Consult a poultry diagnostic laboratory or poultry pathologist for diagnosis and advice on dosage for chickens and turkeys. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

PRECAUTIONS

- Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.
- Use as the sole source of oxytetracycline.
- Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.
- For use in drinking water only.
- Not for use in liquid medicated feeds.

WARNINGS

- Do not feed to birds producing eggs for human consumption.

STORAGE

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F)



Oxytet is a registered trademark of Huvepharma, Inc.
HVP-OXY-PDS-03.18



Oxytet[®] Soluble

(oxytetracycline hcl)

Antibiotic.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE Dosage and indications for use in water proportioners - Add the following amount to 1 gallon of stock solution when proportioner is set to meter at the rate of 1 oz per gallon.

			Oxytet 280g	Oxytet 1400g	Oxytet 1772g
C H I C K E N S	Disease	Treatment Level	Pack/Gallon Stock	Gallons Stock/Pack	Gallons Stock/Pack
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)	20 - 10	25.3 - 12.6
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)	10 - 5	12.6 - 6.3
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)	10 - 5	12.6 - 6.3
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)	20 - 10	25.3 - 12.6
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	½ (140 g)	10	12.6
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption	Varies with age & water consumption	Varies with age & water consumption
S W I N E	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE — Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption	Varies with age & water consumption	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.				

Oxytet 280g (9.8 oz):

This packet contains 102.4g of oxytetracycline hydrochloride and will make:
512 gallons (1936 liters) containing 200 mg oxytetracycline hydrochloride per gallon
128 gallons (484 liters) containing 800 mg oxytetracycline hydrochloride per gallon
This packet will treat 10,240 lbs of swine.
This packet will treat 4,096 lbs of turkey at 25 mg/lb body weight.

Oxytet 1400g (3.09 lbs):

This packet contains 512 g of oxytetracycline hydrochloride and will make:
2560 gallons (9690 liters) containing 200 mg oxytetracycline hydrochloride per gallon
1280 gallons (4845 liters) containing 400 mg oxytetracycline hydrochloride per gallon
640 gallons (2422 liters) containing 800 mg oxytetracycline hydrochloride per gallon
This packet will treat 51,200 lbs of swine.
This packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

Oxytet 1772g (3.91 lbs):

This packet contains 648 g of oxytetracycline hydrochloride and will make:
3240 gallons (12,265 liters) containing 200 mg oxytetracycline hydrochloride per gallon
1620 gallons (6,132 liters) containing 400 mg oxytetracycline hydrochloride per gallon
810 gallons (3,066 liters) containing 800 mg oxytetracycline hydrochloride per gallon
This packet will treat 64,800 lbs of swine.
This packet will treat 25,920 lbs of turkey at 25 mg/lb body weight.

USE DIRECTIONS: Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING

Do not feed to birds producing eggs for human consumption.

RECOMMENDED PACKET

STORAGE CONDITIONS: Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

For Use in Drinking Water Only
Not For Use in Liquid Medicated Feeds
Not For Use in Humans
Keep Out of Reach of Children
HAZARDOUS REMEDY
Restricted Drug (California) - Use only as directed.
NADA 130-435, Approved by FDA