



FOR ANIMAL USE ONLY

INDICATIONS: For use as an aid in prevention and treatment of Ketosis (Acetonemia) in dairy cattle.

DIRECTIONS FOR USE: For use mixed with feed or as a drench.

Prevention: Dairy Cattle: 6-8 fl oz/head/day
Begin 2 weeks before calving and continue for 6 weeks after calving

Treatment: Dairy Cattle: 6-16 fl oz/head/day
Dosage dependant upon severity, to be given for 10 days.

PROPYLENE GLYCOL, U.S.P



For Use as an Aid in Prevention and Treatment of Ketosis

KEEP OUT OF REACH OF CHILDREN

Consult veterinarian if condition does not respond.

LIVESTOCK DRUG

STORAGE: Store at controlled room temperature between 2°-30°C (36°-86°F). Keep container tightly closed, when not in use.

Seller makes no warranty of any kind whether in accordance with directions or not.

ACTIVE INGREDIENT:

Propylene Glycol100%

TAKE TIME



OBSERVE LABEL DIRECTIONS

Manufactured for:
DURVET, INC.

Blue Springs, Missouri 64014

ISSU11XB04

www.durvet.com

NET CONTENTS: ONE GALLON (3.785 L)

Lot No.

Exp. Date

[]

