



# SULFADIMETHOXINE ORAL SOLUTION

ANTIBACTERIAL



## INDICATIONS:

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-Producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - Use for the treatment of shipping fever complex, bacterial pneumonia, calf diphtheria, and foot rot.

## FOR USE ON:

Dairy Calves, Dairy Heifers, Beef Cattle, Broiler and Replacement Chickens, and Meat-Producing Turkeys

## BENEFITS:

- **Broad-spectrum** - An aid in the treatment of coccidiosis, fowl cholera and infectious coryza in chickens; fowl cholera and coccidiosis in turkeys; and shipping fever, bacterial pneumonia, foot rot and calf diphtheria in cattle.
- **Rapid penetration** - Effective blood and tissue levels reached rapidly.
- **Fast-acting** - Works fast because it's not inactivated by minerals in drinking water.
- **Highly palatable** - High palatability assures animals receive the entire dosage.



## ACTIVE INGREDIENT:

Each fluid ounce contains 3.75 gm sulfadimethoxine solubilized with sodium hydroxide.

## USE DIRECTIONS:

Chickens - 0.05% concentration - 25 fluid oz. to 50 gallons of drinking water

Turkeys - 0.025% concentration - 25 fluid oz. to 100 gallons of drinking water

Treatment period: 6 consecutive days.

Dairy Calves, Dairy Heifers, and Beef Cattle - 25mg/lb first day followed by 12.5mg/lb/day for 4 days.

See package label for complete directions.

## PACKAGING:

Gallon, 4 gallons/case    UPC # 7-45801-11957-4



DURVET, INC. ANIMAL HEALTH PRODUCTS  
P.O. Box 279 · Blue Springs, Missouri 64014 · 800/821-5570 · 816/229-9101  
Fax 816/224-3080 · E-mail: info@durvet.com · www.durvet.com