


SpecLinx-50[®]

(Brand of Lincomycin-Spectinomycin soluble powder)
Water Soluble Powder

ANADA 200-380, Approved by FDA



INDICATIONS

For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin.

Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

See reverse side for Administration and Dosage.

BENEFITS

- **Highly soluble:** Mixes easily and completely in water
- **Economical:** A single 375 g packet treats 125 gallons of water
- **Convenient:** Pre-measured packaging provides for handy on-site mixing
- **Stable:** Stores well at room temperature
- **FDA approved**

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1LIN009	75 g pouch	12
1LIN010	375 g pouch	6

SpecLinx-50[®]

(Brand of Lincomycin-Spectinomycin soluble powder)
Water Soluble Powder

ANADA 200-380, Approved by FDA

INGREDIENTS

Each 75 g of SpecLinx-50 contains:

Lincomycin hydrochloride	
Equivalent to lincomycin	16.7 grams
Spectinomycin dihydrochloride pentahydrate	
Equivalent to spectinomycin	33.3 grams
Total Antibiotic Activity	50.0 grams

INDICATIONS

For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin.

Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

DOSAGE

Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.

Administration	
Amount of Spec-Linx 50	Amount of Drinking Water
75 g	25 Gallons
375 g	125 Gallons

For proportioners delivering 1 oz/gallon of drinking water, dissolve the contents of 5 packets of the 75 g pouch or 1 packet of the 375 g pouch size in each gallon of proportioner solution.

IMPORTANT

Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:

Broilers and Layer Replacements (Light and Heavy)

Age (Weeks)	Daily Water intake Gallons/1000	Dosage mg Antibiotic/lb
1	5	50-65

CAUTION

Discard medicated drinking water daily and replace with fresh medicated drinking water. Store at controlled room temperature 20° to 25°C (68° to 77°F).

WARNING

NOT FOR HUMAN USE
KEEP OUT OF REACH OF CHILDREN

HOW SUPPLIED

SpecLinx-50 is supplied in 75 g (2.65 oz) and 375 g (13.2 oz) foil pouches; and 75 g (2.65 oz) plastic jars.

TAKE TIME



OBSERVE LABEL DIRECTIONS