

## SPECTAM<sup>®</sup> Water Soluble Antibiotic

(Spectinomycin)

### Description

SPECTAM is a water soluble concentrate intended as an anti-infective for growing chickens.

Each 1000-gram bottle contains spectinomycin dihydrochloride pentahydrate equivalent to 500 grams of spectinomycin activity.

### Indications


SPECTAM is indicated as an aid in the prevention or control of broiler losses due to chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* (MG) and infectious synovitis associated with *Mycoplasma synoviae* (MS). It is also indicated for use as an effective means of increasing weight gain and improving feed efficiency in floor-raised broilers.

### Benefits

- **Highly soluble**—Mixes readily and thoroughly in water
- **Stable**—Effective over extended periods
- **Versatile**—Promotes feed efficiency and weight gain as well as helping fight disease
- **Easy to use**—Will not clog proportioners
- **FDA approved**

### Packaging

List No.	UNIT PACKAGE
1SPE001	1000-gram, screw-top plastic jar (6 jars per case)



### Dosage and Administration

Mix in drinking water according to dosage directions. This mixture should be the only source of drinking water for the first three (3) days of life, and for one (1) day following each vaccination for CRD and weight gain, and 3-5 days for synovitis.

For CRD associated with MG: Treat water to provide 2 g of spectinomycin activity per gallon.

For infectious synovitis associated with MS: Treat water to provide 1 g of spectinomycin activity per gallon.

For increased weight gain and feed efficiency: Treat water to provide 0.5 g of spectinomycin activity per gallon.

See back and label instructions for complete dosing information.

# Technical Information

---

## SPECTAM<sup>®</sup>

(Spectinomycin Water Soluble Antibiotic)

NADA 038-661, Approved by FDA

### ACTIVE INGREDIENTS

Each gram contains spectinomycin dihydrochloride pentahydrate equivalent to 0.5 grams of spectinomycin activity.

### INDICATIONS

SPECTAM is indicated as an aid in the prevention or control of broiler losses due to chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* (MG) and infectious synovitis associated with *Mycoplasma synoviae* (MS). It is also indicated as an effective means of increasing weight gain and improving feed efficiency in floor-raised broilers.

### DOSAGE & ADMINISTRATION

Mix in drinking water according to dosage directions. This mixture should be the only source of drinking water for the first three (3) days of life, and for one (1) day following each vaccination for CRD and weight gain, and 3-5 days for synovitis.

For CRD associated with MG: Add contents to 250 gallons of water to provide 2 g of spectinomycin activity per gallon. For proportioners, dissolve contents of one-half (1/2) package in one (1) gallon of water. Meter at one (1) oz/gal.

For infectious synovitis associated with MS: Add contents to 500 gallons of water to provide 1 g of spectinomycin activity per gallon. For proportioners, dissolve contents of one-half (1/2) package in two (2) gallons of water. Meter at one (1) oz/gal.

For increased rate of weight gain and improved feed efficiency in floor-raised broiler chickens: Dissolve one-half (1/2) package (500 g) in 500 gallons of water to provide 0.5 g of spectinomycin activity per gallon. For proportioners, dissolve contents of one-half (1/2) package in four (4) gallons of water. Meter at one (1) oz/gal.

### TAKE TIME



### OBSERVE LABEL DIRECTIONS

### WARNING

DO NOT administer this drug within 5 days of slaughter.  
DO NOT administer to laying chickens.

### HOW SUPPLIED

SPECTAM is supplied in 1000 g (35 oz) screw cap, plastic bottles packed 6 per case.



Bimeda, Inc.  
Oakbrook Terrace, IL 60181

SPECTAM is a registered trademark of Bimeda, Inc.  
©2006 Bimeda, Inc. All rights reserved. SPE-TS-0306-3000