

# 117/177mm



## Tylovet<sup>®</sup> Soluble (tylosin tartrate)

For use In chickens, turkeys, swine and honey bees only  
Equivalent to 100 g tylosin base

An Antibiotic  
ANADA# 200-473, Approved by FDA Product 2220



GRANULES FOR ORAL SOLUTION

Chickens, Turkeys, Swine and Honey Bees



3 4800043 17021 24

### INDICATIONS

**Chickens:** As an aid in the treatment of chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* sensitive to tylosin in broiler and replacement chickens. For the control of chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* sensitive to tylosin at time of vaccination or other stress in chickens. For the control of chronic respiratory disease (CRD) associated with *Mycoplasma synoviae* sensitive to tylosin in broiler chickens.

**Turkeys:** For maintaining weight gain and feed efficiency in the presence of infectious sinusitis associated with *Mycoplasma gallisepticum* sensitive to tylosin.

**Swine:** For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae*.

**Honey Bees:** For the control of American Foulbrood (*Paenibacillus larvae*).

### INGREDIENTS

Tylosin (as tylosin tartrate) ..... 100 g

### MIXING DIRECTIONS

**Chickens and Turkeys:** To assure thorough dissolution, place the **Tylovet** (contents of the jar or pouch) in a mixing container and add one gallon of water (3790 mL) to the material. Mix this concentrated solution with water to make 50 gallons (189 liters) of treated drinking water.

**Swine:** To assure thorough dissolution, place the **Tylovet** (contents of the jar or pouch) in a mixing container and add one gallon of water (3790 mL) to the material. Mix this concentrated solution with water to make 400 gallons of treated drinking water resulting in 250 mg/gallon.

**Honey Bees:** Mix 200 mg tylosin in 20 g confectioners/powdered sugar. Use immediately.

**When directed to mix the product with water, always** add the water to the powder. **Do not pour** the powder into the water. Prepare a fresh **Tylovet** solution every three days.

When mixing and handling tylosin, use protective clothing and impervious gloves.

### DIRECTIONS FOR USE

**Chickens** should be treated for three days; however, treatment may be administered for one to five days depending upon severity of infection. Treated chickens must consume enough medicated water to provide 50 mg per pound of body weight per day. Only medicated water should be available to the birds.

**Turkeys** should be treated for three days; however, treatment may be administered for two to five days depending upon severity of infection. Treated turkeys must consume enough medicated water to provide 60 mg per pound of body weight per day. Only medicated water should be available to the birds.

**Swine** should be treated for three to ten days, depending upon severity of infection. Treated swine must consume enough medicated water to provide a therapeutic dose. Only medicated water (250 mg/gal) should be available.

**Honey Bee colonies** should receive three treatments administered as a dust in confectioners/powdered sugar. The 200 mg dose is applied (dusted) over the top bars of the brood chamber once weekly for 3 weeks.

**Notice:** Organisms vary in their degree of susceptibility to any chemotherapeutic. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reconfirmed.

### NOT FOR HUMAN USE

**RESIDUE WARNING:** Chickens must not be slaughtered for food within 24 hours after treatment. Turkeys must not be slaughtered for food within five days after treatment. Swine must not be slaughtered for food within 48 hours after treatment. Do not use in layers producing eggs for human consumption. Honey Bees: The drug should be fed early in the spring or fall and consumed by the bees before the main honey flow begins, to avoid contamination of production honey. Complete treatments at least 4 weeks prior to main honey flow.

**Avoid contact with human skin. Exposure to tylosin may cause a rash.**

### How supplied:

Tylovet<sup>®</sup> Soluble is supplied in pouches of laminated aluminum foil, 100 g of tylosin activity per pouch; or HDPE jars with PP cap with heat sealable coating, 100 g of tylosin activity per jar.

### Storage:

**Store at Room Temperature, 25°C (77°F).**

**(Excursions Permitted to 40°C/104°F). Avoid moisture.**

**Restricted Drug (California) Use Only as Directed.**

**ANADA # 200-473, Approved by FDA**

**To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.**

**Manufactured for:** Huvepharma Inc. Peachtree City, GA 30269, Made in Bulgaria

Lot: 0000000000

Exp. Date: 00.0000

Code: LB444v1-Tyl100-USA1010



C/PP/ALU/LOPE