

# MOTOMCO

# FARM & HOME

**Kills Warfarin  
Resistant Norway  
Rats & House Mice**

**For Indoor and  
Outdoor Rat and  
Mouse Control**

**Ready to Use**

**Pelleted Bait**

**With Brodifacoum**

# JAGUAR<sup>®</sup>



**Rodenticide**

# PLACE PACS

For Use In and Around Agricultural Buildings Only



## KILLS RATS & MICE

ACTIVE INGREDIENT:

Brodifacoum (CAS #56073-10-0): . . . . . 0.005%

OTHER INGREDIENTS\*: . . . . . 99.995%

\*Contains Denatonium Benzoate TOTAL 100.000%

**Net Weight:**

**8 lbs (3.6 kg)**

**73 x 1.76 oz (50 g) Place Pacs**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panels for First Aid and additional precautionary statements.

**FIRST AID****HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE**

<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.</li> <li>Have a person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>
<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.</li> </ul>

**NOTE TO PHYSICIAN**

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Vitamin K<sub>1</sub> is antidotal at doses of 10-20 mg (not mg/kg). Repeat as necessary based on monitoring of prothrombin times.

**TREATMENT FOR PET POISONING**

If animal eats bait, call veterinarian at once.

**NOTE TO VETERINARIAN**

**Anticoagulant Brodifacoum:** For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub> at doses of 2-5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

**USE RESTRICTIONS:** For control only of Norway rats, roof rats, and house mice. For use in and around agricultural buildings only. **Do not apply further than 50 feet from**

**MOTOMCO FARM & HOME**

# JAGUAR<sup>®</sup>

## Rodenticide

# PLACE PACS


**KILLS RATS & MICE****KILLS WARFARIN RESISTANT NORWAY RATS AND HOUSE MICE**

Norway rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT: Brodifacoum (CAS #56073-10-0): . . . . . 0.005%

OTHER INGREDIENTS: . . . . . 99.995%

\*Contains Denatonium Benzoate . . . . . TOTAL 100.000%

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

See side panels for First Aid and additional precautionary statements.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. **[Plastic:]** Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed. **[Paper:]** Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration. In most states, burning is not allowed.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**Net Weight: 8 lbs (3.6 kg)**  
**73 x 1.76 oz (50 g) Place Pacs**

EPA REG. NO.: 12455-94-3240

EPA EST. NO.: 12455-WI-1

Available Exclusively From:



3699 Kinsman Blvd.  
Madison, WI 53704 U.S.A.  
www.motomco.com  
**MADE IN USA**

(Continued from other panel)

**agricultural buildings.** Do not use in homes or other human residences. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. Do not broadcast bait. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 8 pounds of bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats or mice or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:**

**RATS:** Apply 2 to 8 pacs (at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until there no longer are signs of fresh rat activity.

**HOUSE MICE:** Open pac and apply 1/4 to 1/2 oz. of bait per placement in infested areas. Space placements at 8 to 12 foot intervals. Larger placements, up to one unopened pac, may be placed at points of extremely high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Keep away from children, domestic animals and pets. Do not get in eyes or on clothing. All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**User Safety Requirements**

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

#### NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Vitamin K<sub>1</sub> is antidotal at doses of 10-20 mg (not mg/kg). Repeat as necessary based on monitoring of prothrombin times.

#### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once.

#### NOTE TO VETERINARIAN

**Anticoagulant Brodifacoum:** For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub> at doses of 2-5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

#### SEE LABEL ON OUTER PACKAGING FOR ADDITIONAL PRECAUTIONARY INFORMATION

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See outer package label for complete DIRECTIONS FOR USE including USE RESTRICTIONS and APPLICATION DIRECTIONS. It is illegal to sell this product unless it is accompanied by a completed set of its accepted labeling.

**MOTOMCO FARM & HOME**

**JAGUAR<sup>®</sup>**  
**Rodenticide**  
**PLACE PAC**



For Use In and Around Agricultural Buildings Only

**KILLS RATS & MICE**

Norway rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

#### ACTIVE INGREDIENT:

Brodifacoum (CAS #56073-10-0): . . . . . 0.005%

OTHER INGREDIENTS\*: . . . . . 99.995%

\*Contains Denatonium Benzate TOTAL 100.000%

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

SEE LABEL ON OUTER PACKAGING FOR  
ADDITIONAL PRECAUTIONARY INFORMATION

See back panels for First Aid and additional precautionary statements.

**Net Weight: 1.8 OZ (50 g)**

EPA REG. NO. 12455-94-3240

EPA EST. NO. 12455-WI-1

Available Exclusively From:

**MOTOMCO**

3699 Kinsman Blvd.  
Madison, WI 53704 U.S.A.  
www.motomco.com

#### FIRST AID

**HAVE LABEL WITH YOU WHEN  
OBTAINING TREATMENT ADVICE**

#### If Swallowed

- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

#### If in Eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.

**INDIVIDUAL  
SALE IS  
PROHIBITED  
BY LAW**