

GentaMed™

Soluble Powder

(Gentamicin Sulfate)

**ANTIBACTERIAL
FOR SWINE DRINKING WATER ONLY**

INDICATIONS

GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Treponema hyodysenteriae*.

EACH g CONTAINS: Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

BENEFITS

- **Effective:** Stays in the GI tract to fight infection
- **Versatile:** Approved to both control and treat swine Colibacillosis and Dysentery
- **Convenient:** Treats large numbers of swine through the drinking water
- **Economical:** One scoop medicates 240 gallons of drinking water to treat and control colibacillosis at a concentration of the 25 mg/gal
- **Easy to use:** Packaged in screw-top, moisture resistant jars for easy opening and closing



PACKAGING

| LIST NO. | UNIT PACKAGE | CASE SIZE |
|----------|---------------|-----------|
| 1GEN009 | 360 g per jar | 2 |

RESIDUE WARNING

For use in swine drinking water only. Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Bimeda, Inc.

One Tower Lane, Suite 2250
Oakbrook Terrace, IL 60181 USA
Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035
Email: sales@bimedaus.com

See reverse side for Administration and Dosage.



GentaMed™

Soluble Powder

(Gentamicin Sulfate)

ANTIBACTERIAL

FOR SWINE DRINKING WATER ONLY

ANADA 200-494, Approved by FDA

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

DESCRIPTION:

Each gram of GentaMed™ Soluble Powder contains gentamicin sulfate equivalent to 333.33 mg gentamicin base.

INDICATIONS:

GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weaning swine caused by strains of *E. coli* sensitive to gentamicin and for the control and treatment of swine dysentery associated with *Treponema hyodysenteriae*.

DOSAGE AND ADMINISTRATION:

GentaMed™ Soluble Powder is available in a 360 gram jar. The 360 gram jar contains a scoop that provides approximately 18 grams of product when level full.

For Control and treatment of Colibacillosis

Administer GentaMed™ Soluble Powder in drinking water at the recommended level of 25 mg per gallon (1 scoop/240 gallons) for three consecutive days.

For Control and treatment of Swine Dysentery

Administer GentaMed™ Soluble Powder in drinking water at the recommended level of 50 mg per gallon (1 scoop/120 gallons) for three consecutive days.

The concentration of the medication should be adjusted in extremely hot or cold weather to ensure a gentamicin dosage of approximately 0.5 mg/lb/day for 3 days for colibacillosis or 1.0 mg/lb/day for 3 days for swine dysentery. A 50-lb pig will, under average conditions, consume 0.9 gallons of water per day.

For swine dysentery, if the condition recurs, the medication may be repeated. Because of the tendency for the disease to recur on a premise with a history of swine dysentery, a control program should be used following treatment.

DIRECTIONS FOR USE:

To prepare medicated water, mix GentaMed™ Soluble Powder and drinking water according to the following tables:

Proportioner Use

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed™ Soluble Powder to 2 gallons of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 25 mg/gallon.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed™ Soluble Powder to 1 gallon of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 50 mg/gallon.

Bulk Preparation:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed™ Soluble Powder to 240 gallons of water.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed™ Soluble Powder to 120 gallons of water.

Medicated drinking water should not be stored or offered in rusty containers since the drug is quickly destroyed in such containers.

Medicated water should be prepared daily and be the sole source of drinking water for 3 consecutive days.

The daily water consumption figures shown below are approximations for mild temperatures and are presented as a guide only. Actual water consumption varies with environmental temperature, humidity, and diet.

Approximate Daily Water Consumption

| Weight of Pig (lb) | Gallons for 5 Pigs | Gallons for 10 Pigs |
|--------------------|--------------------|---------------------|
| 25 | 3.0 | 6.0 |
| 50 | 4.5 | 9.0 |
| 75 | 6.0 | 12.0 |
| 100 | 7.5 | 15.0 |

SIDE EFFECTS:

There have been no reported side effects following the recommended use of GentaMed™ Soluble Powder in swine.

CONTRAINDICATIONS:

There are no known contraindications to this drug when used as directed.

PRECAUTIONS:

Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

HOW SUPPLIED:

GentaMed™ Soluble Powder is available in a 360 gram jar with 2 jars per shipper.

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com
8GEN001

TAKE TIME



OBSERVE LABEL DIRECTIONS