



Havoc[®]

rodenticide bait



Item No.	Description
116395	Havoc Rodenticide Bait 25 lb. Bag
116398	Havoc Rodenticide Bait 25 lb. Pail
116396	Havoc Rodenticide Bait 50 lb. Bag

Active Ingredient: Brodifacoum

- Acute, single-dose rodenticide containing the active ingredient, 0.005% Brodifacoum
- Kills House Mice, Norway Rats and Roof Rats
- Rodents may consume a lethal dose in a single-night's feeding, with the first dead rodents appearing four or five days after feeding begins.
- Very attractive for rodent control when there is a lot of feed sources around.
- Only 1 to 2 tablespoon amounts will control rodents in an 8 x 12 foot space.
- May be used in and around buildings and structures and inside burrows.
- Use as a supplement to Neogen's current Rotational Baiting Strategy- during periods of heavy rodent infestation.

Product Function:

Havoc Rodenticide Bait contains oats and other food material to make bait very attractive when there is a lot of feed sources around. Rodenticide baits, while the least weatherable bait form, gets superior rodent acceptance because they contain food-grade ingredients, seeds and grains which give rodents a variety of textures and flavors.

Kills Warfarin-Resistant Norway Rats and House Mice. Rodents may consume a lethal dose in a single-night's feeding, with the first dead rodents appearing four or five days after feeding begins.

Application Directions:

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

Norway and Roof Rats:

Apply 4 to 16 ounces Havoc Rodenticide Bait per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are signs of new feeding by rats.

House Mice:

Apply ¼ to ½ ounce (1-2 level tablespoons) of Havoc Rodenticide Bait per placement. Space placements at 8 - 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are signs of new feeding by house mice.

Follow-up:

Replace contaminated or spoiled bait immediately. Using waterproof gloves collect and dispose of all dead, exposed animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.



800-234-5333 (USA/Canada) or 517-372-9200
inform@neogen.com | www.neogen.com

