

Quat-A-Mone™ Sanitizer - Virucide* - Cleaner

ACTIVE INGREDIENTS:

Alkyl (60% C14, 30% C16, 5% C12, 5% C18)
Dimethyl Benzyl Ammonium Chloride 10.0%
Alkyl (68% C12, 32% C14)

Dimethyl Ethylbenzyl Ammonium Chloride..... 10.0%

OTHER INGREDIENTS:..... 80.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See side panel for additional precautionary statements.

FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Est. No. 67589-NC-1

NET CONTENTS: 55 GALLONS

EPA Reg. No. 10324-94-91014

- Cleans, disinfects and deodorizes hard, non-porous surfaces by killing many odor-causing microorganisms.
- Kills Influenza A Virus.
- Is a phosphate-free formulation designed to provide effective cleaning, deodorizing and disinfection in areas where housekeeping is of prime importance in controlling the hazard of cross-contamination on treated surfaces.
- Is a versatile disinfectant & sanitizer for veterinarian, veterinary practice, animal care, animal laboratory, and agricultural and farm premise applications.
- Is for use in federally inspected meat and poultry plants on all hard, non-porous surfaces in inedible product processing areas, non-processing areas and/or exterior areas, federally inspected meat and poultry plants as a floor and wall cleaner for use in all departments, and federally inspected meat and poultry plants as a disinfectant agent for use in all departments.
- This product at 0.5 oz. of this product per 4 gal. of water, this sanitizer fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 2011 Recommendations of the U.S. Public Health Services in waters up to 500 ppm of hardness calculated as CaCO3 when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.
- Can be applied through foaming apparatus and low-pressure sprayers systems. Follow manufacturers' instructions when using this equipment.
- Clear formula.
- Contains no fragrances.
- Contains no phosphorus.
- Cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine or poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities.

For use in: animal research centers, animal quarantine areas, animal holding areas, farms, animal breeding facilities, pet animal quarters, zoos, tack shops, operating rooms, washing areas, waiting rooms, examination rooms and other animal care facilities, dairy, sheds, pens and stalls, swine quarters, equine quarters, brooder houses, seed houses and veal, calving, hog, cattle and horse operations, farm vehicles, federally inspected meat and poultry plants, food handling and processing areas, USDA inspected food-processing facilities, federally inspected meat and poultry plants, egg processing plants.

For use on: poultry/turkey equipment, egg receiving and egg holding areas, meat packing plant surfaces such as livestock vehicles and holding pens.

DISINFECTION PERFORMANCE: This product kills the following bacteria in 10 minutes at 2 oz. per 5 gal. of water and 5% soil on hard, non-porous surfaces: *Botrytis cinerea*, *Burkholderia cepacia*, *Campylobacter jejuni*, *Corynebacterium ammoniogenes*, *Enterococcus faecium* Vancomycin Resistant, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella enterica*, *Salmonella typhi*, *Shigella sonnei*, *Staphylococcus aureus*, *Staphylococcus aureus* (MRSA), *Staphylococcus aureus* (CA MRSA), *Yersinia enterocolitica*.

VRUCIDAL* PERFORMANCE: This product kills the following viruses in 10 minutes at 2 oz. per 5 gal. of water and 5% soil on hard, non-porous surfaces: Avian Influenza A (H5N1) Virus, Avian Influenza A/Turkey/Wisconsin Virus, Herpes Simplex Type 1 Virus, Herpes Simplex Type 2 Virus, Human Coronavirus, Influenza A (H1N1) Virus, Influenza A2/Japan Virus, Vaccinia Virus. †Indicates a 2-minute contact time is required for this claim.

ANIMAL PREMISE VRUCIDAL* PERFORMANCE: This product kills the following viruses in 10 minutes at 2 oz. per 5 gal. of water and 5% soil on hard, non-porous surfaces: Avian Influenza A (H5N1) Virus, Avian influenza A/Turkey/Wisconsin Virus, Avian Reovirus, Bovine Viral Diarrhea Virus, Canine Coronavirus, Canine Distemper Virus, Equine Arteritis Virus, Infectious Bovine Rhinotracheitis Virus, Infectious Bronchitis Virus, Infectious Laryngotracheitis Virus, Newcastle Disease Virus, Porcine Respiratory & Reproductive Virus, Porcine Rotavirus, Pseudorabies Virus, Transmissible Gastroenteritis Virus.

FOOD CONTACT SURFACE SANITIZING PERFORMANCE: This product is an effective food contact sanitizer in 1 minute at 0.5 oz. per 4 gal. of 500 ppm hard water on hard, non-porous surfaces: *Aeromonas hydrophila*, *Campylobacter jejuni*, *Clostridium perfringens*, *Enterobacter sakazakii*, *Enterococcus faecalis* Vancomycin Resistant, *Escherichia coli* O111:H8, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Salmonella typhi*, *Shigella dysenteriae*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Yersinia enterocolitica*.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Rinse hard, non-porous food contact surfaces with potable water after application of product.

Before using this product, food products and packaging materials must be removed from the room or carefully protected.

FOR USE AS A GENERAL DISINFECTANT, VIRUCIDE*, DEODORIZER, CLEANER: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 2 oz. of this product per 5 gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain wet for 10 minutes. 4. Allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. 5. Prepare a fresh solution daily or when visibly dirty.

FOR USE AS A DISINFECTANT/CLEANER IN FOOD PROCESSING PLANTS/PREMISES, FOOD STORAGE AREAS AND EQUIPMENT: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 1 oz. of this product per gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain wet for 10 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. When disinfecting food contact surfaces used for food preparation, rinse surfaces thoroughly

with potable water. This product must not be used to clean the following surfaces: utensils, glassware, dishes or interior surfaces of processing equipment. **Notes:** All food contact surfaces such as appliances and kitchen countertops must be rinsed with potable water after treatment. Do not use this product to clean or disinfect glassware, utensils, dishes or interior surfaces of appliances.

FOR USE AS A SANITIZER AFTER DISINFECTION IN FOOD PROCESSING PLANTS/PREMISES, RENDERING AND DRESSING PLANTS, FOOD STORAGE AREAS AND EQUIPMENT: 1. Pre-clean heavily soiled areas. 2. Apply use solution of 2 oz. of this product per 5 gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. 3. Treated surfaces must remain wet for 10 minutes. 4. Wipe dry with a clean cloth or allow to air dry. 5. Thoroughly wash equipment and utensils with a compatible detergent. 6. Rinse with potable water. 7. Sanitize in a solution of 0.5-1 oz. of this product per 4 gal. of water (200-400 ppm active quat) (or equivalent use dilution). Surfaces must remain wet for 1 minute. 8. Drain or allow to air dry. 9. Prepare a fresh solution daily or when visibly dirty.

ANIMAL PREMISES: Prior to use of this product, remove all animals and feeds from premises, animal transportation vehicles, and enclosures. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIRUCIDE*: Prepare a use solution of 2 oz. of this product per 5 gal. of water. Apply use solution to disinfect hard, non-porous surfaces with a sponge, brush, cloth, by immersion, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. Treated surfaces must remain wet for 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

HATCHERIES: Use to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard, non-porous surfaces. Use 2 oz. of this product per 5 gal. of water. Leave all treated surfaces wet for 10 minutes or more. Allow to air dry.

VEHICLES: To clean hard, non-porous surfaces on vehicles including mats, crates, cabs, and wheels, use 2 oz. of this product per 5 gal. of water. Apply use solution to hard, non-porous surfaces on vehicles. Leave treated surfaces wet for 10 minutes. Allow to air dry.

FOGGING IN FOOD PREMISES

ALL SURFACES MUST BE CLEANED AND DISINFECTED IN ACCORDANCE WITH LABEL DIRECTIONS PRIOR TO FOGGING. DIRECTIONS FOR FOGGING IN DAIRIES, BEVERAGE AND FOOD PROCESSING PLANTS: Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After disinfecting, fog desired areas using 0.5 quart per 1,000 cubic feet of room area with a solution containing 0.8 oz. of product per gal. of water. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. When fogging is complete, ventilate buildings and other closed spaces. All food contact surfaces must be sanitized with an EPA approved food contact sanitizer solution prior to use. All food contact surfaces must be thoroughly rinsed with potable water prior to sanitizing. **Note:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH, goggles, long sleeves, gloves and long pants.

FOGGING IN POULTRY HOUSES

ALL SURFACES MUST BE CLEANED AND DISINFECTED IN ACCORDANCE WITH LABEL DIRECTIONS PRIOR TO FOGGING. TREATMENT OF HATCHERY ROOMS USING FOGGING DEVICES: Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.

Close room off so fog is confined to room to be treated. Mix 28 oz. of this product per gal. of water. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. When fogging is completed ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed or dried. Thoroughly clean all treated surfaces with soap or detergent and rinse with water. **Note:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH, goggles, gloves, long sleeves and long pants.

TREATMENT OF INCUBATORS AND HATCHERS USING FOGGING DEVICES: Only for treatment of setters and hatchers after poultry/chicks/eggs have been removed. Not for treatment of hatchers which contain chicks/eggs. Mix 7.2 oz. of this product per gal. of water. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Fog 3-8 oz. of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatchery treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog. It is acceptable to fog setters and hatchers with a 0.55 oz. of this product per gal. of water solution on an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. When fogging is complete, ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers, and other treated equipment with soap or detergent, and rinse with potable water before reuse. **Note:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH, goggles, gloves, long sleeves and long pants.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, chemical-resistant gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, oysters, and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

G-18

Sold By: ClearView Enterprises, LLC
451 Agnes Drive
Tontitown, AR 72770



Clear View Enterprises