PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes eye damage and skin damage. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL AND CHEMICAL HAZARDS: Do not use, pour, spill, or store near heat or open flame.

Concentrated broad spectrum iodophor for use as a one-step cleaner/disinfectant and no-rinse sanitizer in:

- Poultry and Livestock Drinking Water
- Dairy and Poultry Farms Meat and Poultry Plants
- Sanitizing Commercial Eggs
- Egg Processing Plants
- Food Processing Plants

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For disinfection, remove gross filth or heavy soil. Apply lodis with a cloth, mop, or mechanical coarse spray device. When applied with a mechanical coarse spray device, surfaces must be spraved until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled.

Disinfection: Veterinary Clinics, Fish Hatcheries, Schools, Farm Premises, and Poultry Houses:

Remove gross filth or heavy soil. Apply 1/2 fl. oz. lodis per gal. of water (or equivalent use dilution) to walls, floors, and other hard, non-porous surfaces such as tables, chairs, countertops, tile, and porcelain surfaces. Allow surfaces to remain wet for at least 10 minutes. For heavily soiled areas, a preliminary cleaning is required.

For farms and poultry premise use, prior to disinfection, all poultry, other animals, and their feeds must be removed from the premises. This includes emptying all troughs, racks, and other feeding and watering appliances. Remove all litter and droppings from floors, walls, and other surfaces occupied or traversed by poultry or other animals.

After 10 minutes disinfection, ventilate buildings, coops, and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set. or dried.

SANITIZING FOOD PROCESSING EQUIPMENT AND UTENSILS:

Remove heavy soils by flushing, scraping, or mopping. Wash with 1 fl. oz. lodis per gal. of water (or other good cleaner). Rinse with clean water and sanitize surfaces with 1 fl. oz. lodis per 5 gal, of water (or equivalent use dilution). Surfaces should remain in contact with the use solution for 1-2 minutes. Drain well and air dry. No potable water rinse is required.

SANITIZING FOOD GRADE EGG SHELLS:

To sanitize clean shell eques intended for food or food products, spray with a solution of 1 fl. oz. of lodis in 5 gal. of water (or equivalent use dilution). The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been freshly washed may be sanitized with this compound only if the eggs are rinsed prior to application of the compound. A subsequent potable water rinse is not required. Eggs must be reasonably dry before casing or breaking. The solution must not be reused for sanitizing eggs.

SANITIZING POULTRY AND LIVESTOCK DRINKING WATER:

Add 1 fl. oz. of lodis per 10 gal. of drinking water.

SHOE BATH SANITIZER:

To prevent tracking harmful organisms into animal areas, scrape shoes and place in a 3 fl. oz. lodis per gal. of water solution (or equivalent use dilution) for 30 seconds prior to entering area

DEODORIZING:

lodis destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab surfaces to be treated with a solution containing 1/2 fl. oz. lodis per gal. of water (or equivalent use dilution). Allow treated surfaces to drain and air dry

IODIS

Disinfectant-Cleaner-Sanitizer-Deodorizer

With 5% Organic Soil and 400 ppm Hard Water Tolerance For Hospital, Institutional, Industrial, School, Agriculture, and Dairy

Active Ingredient:	
lodine*	
Inert Ingredients:	
Total	
* from Nonylphenoxypoly (ethyleneoxy) ethanol-iodine complex	
EPA Reg. No. 66171-102	EPA Est. No. 66171-CA-001

66171-TN-00⁻ 61282-WI-01 47000-IA-0 (First letter of batch number indicates establishment

KEEP OUT OF REACH OF CHILDREN DANGER

See side panels for additional Precautionary Statements

FIRST AID	
If in Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center, or doctor, or 1-800-498-5743 immediate! for treatment advice.
If on Skin or on Clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center, doctor, or 1-800-498-5743 immediately fo treatment advice.
lf Inhaled:	Move person to fresh air. If not breathing, call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth, if possible. Call a poison control center, doctor, or 1-800-498-5743 for treatmen advice.
If Swallowed:	Call a poison control center, doctor, or 1-800-498-5743 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control cente or doctor. Do not give anything by mouth to an unconscious person.
	HOT LINE NUMBER

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Store away from heat. Do not transfer to another container. Do not reuse empty container. PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for quidance.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or offer for reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

GENERAL: Consult Federal, State, or local disposal authorities for approved alternative procedures

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitations of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

To the extent permitted by applicable law, Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use. To the extent permitted by applicable law. MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE. NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY, EXCEPT AS STATED ABOVE.

To the extent permitted by applicable law, in no event shall Manufacturer or Seller be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES, OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY, OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Manufacturer.

For non-emergency (e.g., current product information), call: 1-800-621-8829.



Manufactured by: Preserve International 944 Nandino Blvd., Lexington, KY 40511 U.S.A. Preserve International is a wholly-owned subsidiary of Neogen Corporation. L5572-1117

系 NEOGEN



Net Contents: 55 gal. (208 L)