

Farm Animal Care

Disinfectant Cleaner & Deodorizer

Ready to Use Wipes



EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1 | EPA Est. No. 75272-WI-1
See EPA Est. No. on product container

Manufactured by:

Virox Technologies Inc.

2770 Coventry Road, Oakville, ON L6H 6R1
1-800-387-7578 | www.virox.com

**KEEP OUT OF
REACH OF CHILDREN**

This product is a one-step³ disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile and stainless steel associated with high touch surfaces, animal equipment, animal crates, automatic feeders^A, buckets, brushes, exterior surfaces of watering appliances, farm vehicles, feed racks^A, foot baths, grooming equipment, handling equipment, pens, transportation crates, -or- any hard, non-porous washable surface where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Claims

*VIRUCIDAL in 1 minute unless otherwise noted:

Broad-spectrum *Virucidal activity. Kills Avian Influenza A Virus on pre-cleaned environmental hard, non-porous surfaces.

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Avian Reovirus
Bovine Viral Diarrhea virus
Cytomegalovirus
Duck Hepatitis B virus
Equine Herpes virus type 1
Equine Influenza A Virus (H3N8) (EIV)
Foot and Mouth Disease virus (FMDv) – 3 minutes
Hepatitis B Virus (HBV) – 30 seconds
Hepatitis C Virus (HCV) – 30 seconds
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B – 30 seconds
Human Coronavirus
Influenza A Virus
Influenza B Virus
Norovirus
Marek's Disease virus
Newcastle disease virus
Pandemic Influenza A Virus H1N1
Parainfluenza Virus Type 3
Porcine Epidemic Diarrhea virus (PEDv)
Porcine Respiratory & Reproductive Syndrome virus (PRRS)
Porcine Rotavirus
Pseudorabies virus
Respiratory Syncytial Virus (RSV)
Rhinovirus Type-37
Rotavirus (WA strain)

SARS-Related Coronavirus 2 (Human Coronavirus SARS-CoV-2 (the virus that causes COVID-19²))

Vaccinia Virus

Vesicular Stomatitis Virus

Meets surface disinfection recommendations under the OSHA Bloodborne Pathogen Standard for *HBV, *HCV & *HIV-1

BACTERICIDAL in 1 minute:

Acinetobacter baumannii (Multi-Drug Resistant)
Bordetella bronchiseptica
Bordetella pertussis
Burkholderia cepacia
Enterobacter aerogenes
Enterobacter cloacae (New Delhi metallo-β-lactamase 1 (NDM-1))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Escherichia coli
Escherichia coli O157:H7
Escherichia coli (Extended spectrum β-lactamase (ESBL))
Escherichia coli (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended spectrum β-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Legionella pneumoniae
Listeria monocytogenes
Pasteurella multocida
Pseudomonas aeruginosa
Pseudomonas aeruginosa – (Metallo-beta-lactamase (MBL) positive)
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus Community Acquired Methicillin Resistant Genotype USA 400 (CA-MRSA 400)
Staphylococcus aureus (Methicillin Resistant (MRSA))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Staphylococcus pseudintermedius
Stenotrophomonas (Pseudomonas) maltophilia
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Streptococcus pyogenes

FUNGICIDAL in 30 seconds, unless otherwise noted:

Candida albicans – 1 minute
Microsporum canis – 1 minute
Trichophyton interdigitale

TUBERCULOCIDAL in 1 minute:

Mycobacterium bovis (BCG)

NON-FOOD CONTACT SANITIZER in 10 seconds against:

Acinetobacter baumannii
Campylobacter jejuni
Enterobacter aerogenes
(Vancomycin Resistant) *Enterococcus faecalis* (VRE)
Escherichia coli O157:H7
Listeria monocytogenes
Pseudomonas aeruginosa
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Staphylococcus aureus
(Methicillin Resistant) *Staphylococcus aureus* (MRSA)

Mold/Mildew (Mildewcidal Activity) – at a 4 minute contact time:

Aspergillus brasiliensis (formerly known as *Aspergillus niger*)

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. A potable water rinse is required for food contact surfaces.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

For Use as One-Step Cleaner/Disinfectant

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous surfaces.
3. Allow surface to remain visibly wet for 1 minute.[†]
4. Allow to air dry.

For Use as a Cleaner and/or Deodorizer

1. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
2. Allow to air dry or rinse.

For Use in Animal Housing Facilities such as Barn and Farm Premises

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including pens, stalls, chutes, runs and barns occupied by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from canister and wipe hard, non-porous surfaces.
5. All surfaces must remain visibly wet for 1 minute[†].
6. Allow to air dry or rinse with potable water if necessary.
7. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Mold and Mildew

1. Pre-clean surfaces.
2. Pull towelette from canister and wipe product onto hard, non-porous surfaces until completely wet.
3. Allow the surface to remain visibly wet for 4 minutes.
4. Allow to air dry.

For Use as a Non-Food Contact Sanitizer

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain visibly wet for 10 seconds.
4. Allow to air dry.

To Kill *Trichophyton interdigitale*:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surfaces to remain visibly wet for 30 seconds.
4. Allow to air dry.

For Use as a Tuberculocide:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain visibly wet for 1 minute.
4. Allow to air dry.

To Kill Foot and Mouth Disease virus [FMDv]:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surfaces to remain visibly wet for 3 minutes.
4. Allow to air dry

***KILLS HIV-1 AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of (inanimate) hard, non-porous surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain visibly wet for 30 seconds to kill HIV-1, HBV and HCV.

Disposal of Infectious Material: Cleaning materials used that may contain blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL:

PESTICIDE STORAGE: Store this product in a cool dry area, away from direct sunlight and heat. Do not contaminate water, food or feed by storage and disposal.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture. Non-refillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

USER SHOULD READ S.D.S. BEFORE USING PRODUCT

SDS is available online at www.virox.com

DISCLAIMER: You must read and follow the instructions on the label for our product and use the product for the particular applications specified on the product label. To the full extent permitted under applicable law, Virox Technologies Inc. shall not be liable for any damages whatsoever, arising out of or otherwise in connection with any failure to follow those instructions or any use of its products which is not specified on the product's label. The instructions on the product label are not a warranty, express or implied, and Virox Technologies, Inc. expressly disclaims any implied warranties, including any warranty of fitness for a particular purpose. Under applicable law, some of the above waivers or disclaimers of warranty, or exclusions of liability for certain types of damages may not apply to you. In such cases, our liability will be limited to the fullest extent permitted by applicable law.

²Accelerated Hydrogen Peroxide® technology and designs are trademarks of Diversey Inc. "Accelerated" does not imply increased product efficacy.

^ΔRinse with potable water after disinfecting.

[†]Use a 3 minute contact time against Foot and Mouth Disease virus [FMDv].

^{*}Hard, non-porous surfaces

³When used according to disinfection use directions.